arXiv:1010.5104v2 [physics.gen-ph] 26 Oct 2010

Viscose Cosmology in $f(R)$ Gravity for Kasner Type Metric

Kh. Saaidi[1], **A. Aghamohammadi**[2], **H. Hossienkhani**[3]
Department of Physics, Faculty of Science, University of Kurdistan,
Sanandaj, Iran

Abstract

We study $f(R)$ modifications of Einstein's gravity in a viscous Bianchi type I for a Kasner form metric. We get the $f(R)$ function that satisfy dominant energy condition.

1 Introductions

Observation data[1, 2, 3] indicates that there are several shortcoming of standard gravity (SGR)[4, 5, 6] related to cosmology, large scale structure and quantum field theory. Several attempts preformed to solve these problems by adding new and unjustified components in order to give adaptable picture with present universe. Almost all of these considerations are mainly investigated in a spatially flat homogeneous and isotropic universe described by Friedmann-Robertson-Walker(FRW) metric. The theoretical studies and experimental data, which support the existence of anisotropic phase, leads to consider the model of universe with anisotropic background. Since, the universe is almost isotropic at large scale, the studying of the possible effects of anisotropic universe in the early time makes the Bianchi I type (BI) model as a prime alternative for studding. And solving Einstein equation for a BI universe in the presence of dust, stiff matter and cosmological constant done in [7, 8, 9]. A kind of approach to be studied in the present work is allow an anisotropy in the early universe, and explore where this assumption leads. Although, concerning with present universe, it is known that the anisotropy is small. Thus Bunnet *et al.*[10] have recently done a statistical analysis of 4-yr data from the COBE satellite, and concluded that the present amount of shear $(\sigma/H)_0$ is lass than 3×10^{-9}. In the the early universe, however, the anisotropy may have been larger. The possibility of a state of anisotropy at

---

[1] ksaaidi@uok.ac.ir
[2] agha35484@yahoo.com
[3] Hossienhossienkhani@yahoo.com

very early times is a very natural to investigate, as an attempt to explain, the large local anisotropic that we observe in the universe today in galaxies and supergalaxies[11]. Some other recent papers dealing with anisotropic cosmologist are [12, 13, 14, 15].
The present paper organized as follows: In sec.2, we have reviewed $f(R)$ models, the conformal transform between Jordan frame and Einstein frame and metric of Kasner form metric. In sec.3, we have considered the field equation of $f(R)$ gravity theory in Kasner form. Sec. 4, is related to considering the set of solutions of the field equations. Finally, the latter section is devoted to conclusions.

2 Preliminary

2.1 $f(R)$Theory

$f(R)$ theory is a modified theory of gravity, in which the Einstein-Hilbert lagrangian i.e. R, is replaced by an arbitrary function of scalar curvature i.e. $f(R)$. So the action for a $f(R)$ theory of gravity takes the following form.

$$S_{f(R)} = \int d^4x\sqrt{-g}\left(\frac{f(R)}{2} + \kappa\mathcal{L}_m(g_{\mu\nu,\psi})\right), \tag{1}$$

where $g_{\mu\nu}$ is the metric of space time, ψ is the matter field and $\mathcal{L}_m(g_{\mu\nu,\psi})$ is the matter lagrangian. Here we choose $\kappa = 1$. We assume that the metric $g_{\mu\nu}$ is only independent variable and by variation of the action with respect $g_{\mu\nu}$ gives:

$$R_{\mu\nu}f'(R) - \frac{1}{2}f(R)g_{\mu\nu} - \nabla_\mu\nabla_\nu f'(R) + g_{\mu\nu}\Box f'(R) = T^m_{\mu\nu}, \tag{2}$$

where $T^m_{\mu\nu} = -\frac{1}{\sqrt{-g}}\frac{\partial(\sqrt{-g}\mathcal{L}_m)}{\partial g^{\mu\nu}}$.

2.2 Transformation to a Scalar-Tensor Theory

As regards, the $f(R)$ theory can be recast as a scalar tensor theory via a famous conformal transformation as

$$e^{-2\beta\phi} = f'(R), \tag{3}$$

we can rewrite the action (1) as follows

$$S_{ST} = \int d^4x\sqrt{-\bar{g}}\left(\frac{\bar{R}}{2} - \frac{1}{2}\bar{g}^{\mu\nu}\nabla_\mu\phi\nabla_\nu\phi - V(\phi) + \mathcal{L}(e^{-2\beta\phi}\overline{g}_{\mu\nu}, \psi_i)\right), \tag{4}$$

where, we define the Einstein frame metric $\bar{g}_{\mu\nu}$ by a conformal transformation $\bar{g}_{\mu\nu} = e^{-2\beta\phi} g_{\mu\nu}$. Also the potential $V(\phi)$ is given by

$$V(\phi) = \frac{Rf'(R) - f(R)}{2f'(R)^2}. \tag{5}$$

Then, variation with respect to the independent variables $\bar{g}_{\mu\nu}$ and ϕ give rise to the following field equations

$$\bar{R}_{\mu\nu} - \frac{1}{2}\bar{R}\bar{g}_{\mu\nu} = \nabla_\mu\phi\nabla_\nu\phi + \bar{g}_{\mu\nu}\left[\frac{1}{2}(\nabla\phi)^2 + V(\phi)\right] + \bar{T}^m_{\mu\nu}, \tag{6}$$

$$\bar{\Box}\phi = V'(\phi) - \beta\bar{T}^m. \tag{7}$$

In the above expression, the covariant derivatives, $\bar{\nabla}_\mu$ obey $\bar{\nabla}_\mu\bar{g}_{\mu\nu} = 0$, and all indices are lowered and raised with $\bar{g}_{\mu\nu}$ unless stated other wise.

2.3 The Kasner Form Metric

The line element of the Bianchi type-class I can be expressed as follow

$$ds^2 = dt^2 - A^2(t)dx^2 - B^2(t)dy^2 - C^2(t)dz^2, \tag{8}$$

where, the metric function, A, B, C, being the function of time, t, only. This model is an anisotropic generalization of the Friedmann model with Euclidean spatial geometry. Where, the expansion factors, A, B, C, determined via Einstein equation. At the moment, we shall restrict ourselves to the Kasner form of the metric as:

$$ds^2 = dt^2 - t^{2p_1}dx^2 - t^{2p_2}dy^2 - t^{2p_3}dz^2, \tag{9}$$

where p_1, p_2, p_3 are three parameters which we will require to be constant. The space is anisotropic, if at least two of the three $p_is (i = 1, 2, 3)$ are different.

3 The Field Equation of $f(R)$ Theory of Gravity in the Kasner Form Metric

By making use of (6) and (7), we can derive the field equation. Considering, the Kasner form metric (9) and introduce the symbols $S \equiv p_1 + p_2 + p_3$ and $Q \equiv p_1^2 + p_2^2 + p_3^2$, the components of Ricci tensor R_{00} and R_{ii} obtain as follows:

$$\bar{R}_{00} = \frac{S - Q}{t^2} + \frac{S\beta\dot{\phi}}{t} + 3\beta\ddot{\phi}, \tag{10}$$

$$\bar{R}_{ii} = t^{2p_i}\left(\frac{p_i(S-1)}{t^2} - \frac{\beta\dot{\phi}}{t}(S+2p_i) + 2\beta^2\dot{\phi}^2 - \beta\ddot{\phi}\right), \tag{11}$$

on the other hands, by considering the energy momentum tensor for a viscous fluid as

$$T_{\mu\nu} = \left(\rho + (P - \xi\theta)\right) u_\mu u_\nu - (P - \xi\theta) g_{\mu\nu} + 2\eta\sigma_{\mu\nu}, \tag{12}$$

where u_μ, ρ, P, ξ and η are the fluid's four velocity, the energy density, the isotropic pressure, the bulk and shear viscosities, respectively. The scalar expansion and traceless shear tensor are

$$\theta_{\mu\nu} = \frac{1}{2}\left(u_{\mu;\alpha}h^\alpha_\nu + u_{\nu;\alpha}h^\alpha_\mu\right), \tag{13}$$

$$\sigma_{\mu\nu} = \theta_{\mu\nu} - \frac{1}{3}h_{\mu\nu}\theta, \tag{14}$$

respectively. Here $h_{\mu\nu} = u_\mu u_\nu - g_{\mu\nu}$ is project tensor. The trace of the (6) gives :

$$\bar{R} = (\bar{\nabla}\phi)^2 + 4V(\phi) - \bar{T}^m, \tag{15}$$

substituting, (15) into (6) gives

$$\bar{R}_{\mu\nu} = e^{2\beta\phi}(T_{\mu\nu} - \frac{1}{2}Tg_{\mu\nu}) + e^{-2\beta\phi}g_{\mu\nu}V(\phi) + \nabla_\mu\phi\nabla_\nu\phi. \tag{16}$$

So, the components $\bar{R}_{00}$, $\bar{R}_{ii}$ are as follow;

$$\bar{R}_{00} = e^{2\beta\phi}\left[\rho - \frac{1}{2}(\rho - 3(P - \xi\theta))\right] + e^{-2\beta\phi}V(\phi) + \dot{\phi}^2, \tag{17}$$

$$\bar{R}_{ii} = \left[\frac{e^{2\beta\phi}}{2}\left(\rho - (P - \xi\theta) + \frac{4\eta}{t}(\frac{S}{3} - p_i)\right) - e^{-2\beta\phi}V(\phi)\right]t^{2p_i}. \tag{18}$$

By using equations (10), (11), (17) and (18) one can arrive at.

$$\begin{aligned} e^{2\beta\phi}\left[\rho - \frac{1}{2}(\rho - 3(P - \xi\theta))\right] &+ e^{-2\beta\phi}V(\phi) + \dot{\phi}^2 \\ &= \frac{S-Q}{t^2} + \frac{S\beta\dot{\phi}}{t} + 3\beta\ddot{\phi}, \end{aligned} \tag{19}$$

$$\begin{aligned} p_i\left(-2\eta t e^{2\beta\phi} + \frac{(1-S)}{t^2} + \frac{2\beta\dot{\phi}}{t}\right) &+ \frac{S\beta\dot{\phi}}{t} - 2\beta^2\dot{\phi}^2 + \beta\ddot{\phi} \\ &= -\frac{e^{2\beta\phi}}{2}\left(\rho - (P - \xi\theta) + \frac{4\eta S}{3t}\right) + e^{-2\beta\phi}V(\phi). \end{aligned} \tag{20}$$

Since the $p_i s (i = 1, 2, 3)$ are all linearly independent, we get for the shear viscosity the following relation:

$$\eta = (\frac{1-S}{2t} + \beta\dot{\phi})e^{-2\beta\phi}. \tag{21}$$

Then by substituting, equation (21) into the right hand side of the the (20) we get

$$e^{2\beta\phi}\left(\rho - (P - \xi\theta)\right) = 4\beta^2\dot{\phi}^2 - 2\beta\ddot{\phi} - \frac{10}{3}\frac{S\beta\dot{\phi}}{t} - \frac{2S(1-S)}{3t^2} + 2e^{-2\beta\phi}V(\phi), \tag{22}$$

combining, (22) and (19), we obtain explicit expression for ρ and p as follows

$$e^{2\beta\phi}(P-\xi\theta) = \frac{1}{2t^2}\left(\frac{4}{3}S - Q - \frac{S^2}{3}\right) + \frac{4}{3}\frac{S\beta}{t}\dot{\phi} + 2\beta\ddot{\phi} - (\beta^2 + \frac{1}{2})\dot{\phi}^2 - e^{-2\beta\phi}V(\phi), \tag{23}$$

$$e^{2\beta\phi}\rho = \frac{1}{2t^2}(S^2 - Q) + e^{-2\beta\phi}V(\phi) - 2\beta\frac{S}{t}\dot{\phi} + (3\beta^2 - \frac{1}{2})\dot{\phi}^2. \tag{24}$$

Substituting (23) and (24) into (7), we get the equation of motion for the scalar field ϕ, as follow;

$$\ddot{\phi} + \dot{\phi}\frac{S}{t} - \beta\dot{\phi}^2 = \frac{e^{-2\beta\phi}}{(1-6\beta^2)}\big(V'(\phi) - 4\beta V(\phi)\big) + \frac{C}{t^2}, \tag{25}$$

where $C = \frac{\beta}{(1-6\beta^2)}\big(2S - S^2 - Q\big)$.

4 The Solutions

In this section we want to solve equation (25). It is seen that this equation has a c/t^2 term (c is a constant), therefore the other terms should phrase from order to $\mathcal{O}(1/t^2)$ term. As a result, we suggest the field ϕ to be as

$$\phi = \phi_0 \ln(\frac{t}{t_0}). \tag{26}$$

By this choosing, we can considering three class solution as follows

4.1 Case I

A class solution of (25) is as

$$\frac{e^{-2\beta\phi}}{(1-6\beta^2)}(V'(\phi)-4\beta V(\phi))=0, \tag{27}$$

$$\ddot{\phi}+\dot{\phi}\frac{S}{t}-\beta\dot{\phi}^2-\frac{C}{t^2}=0. \tag{28}$$

Solving (27), we have

$$V(\phi)=V_0 e^{4\beta\phi}. \tag{29}$$

Then by substituting (26) in (28), we can obtain an equation which $\phi_0^{(I)}$ satisfy it, as

$$(S-1)\phi_0^{(I)}-\beta(\phi_0^{(I)})^2-C=0. \tag{30}$$

Also we can obtain ρ, P and η for this situation as

$$\rho=V_0+\frac{\rho_0}{t^{2(\beta\phi_0^{(I)}+1)}}, \tag{31}$$

$$P=-V_0+\frac{S}{t}\xi+\frac{P_0}{t^{2(\beta\phi_0^{(I)}+1)}}, \tag{32}$$

and

$$\eta=\left(\frac{1-S}{2}+\beta\phi_0^{(I)}\right)\frac{t_0^{2\beta\phi_0^{(I)}}}{t^{(2\beta\phi_0^{(I)}+1)}}. \tag{33}$$

Where

$$\rho_0=\left(\frac{S^2-Q}{2}+\left[(3\beta^2-\frac{1}{2})\phi_0^{(I)}-2\beta S\right]\phi_0^{(I)}\right)t_0^{2\beta\phi_0^{(I)}}$$

$$P_0=\left(\frac{2}{3}S-\frac{Q}{2}-\frac{S^2}{6}+\left(\frac{4}{3}S\beta-2\beta-(\beta^2+\frac{1}{2})\phi_0^{(I)}\right)\phi_0^{(I)}\right)t_0^{2\beta\phi_0^{(I)}}$$

So that by using (29) and (5) one can obtain the available $f(R)$ as

$$f(R)=-2V_0+c_I R, \tag{34}$$

where c_I is the integration constant and $2V_0$ is similar to cosmological constant Λ. So that in this class of solution, the function $f(R)$ is similar to the standard Einstein general relativity. For the case $\beta\phi_0^I \simeq 0$, which is equivalent with $f'(R)=1$, this theory is reduce to standard Einstein general relativity theory. In this case the quantity ρ, P and η are in agreement with [16].

4.2 Case II

The another class of solution is as

$$\begin{aligned} \frac{e^{-2\beta\phi}}{(1-6\beta^2)}(V'(\phi) - 4\beta V(\phi)) &= \frac{B}{t^2}, & (35) \\ \ddot{\phi} + \dot{\phi}\frac{S}{t} - \beta\dot{\phi}^2 - \frac{B+C}{t^2} &= 0. & (36) \end{aligned}$$

Where B is the constant, by solving (35) we have

$$V(t) = V_0 t^{4\beta\phi_0^{(II)}} - k\frac{t^{2\beta\phi_0^{(II)}-2}}{t_0^{2\beta\phi_0^{(II)}}}, \tag{37}$$

where

$$k = \frac{B(1-6\beta^2)\phi_0^{II}}{2(1+\beta\phi_0^{II})}. \tag{38}$$

Then by substituting (26) in (36), we can obtain an equation which $\phi_0^{(II)}$ satisfy it as

$$(S-1)\phi_0^{(II)} - \beta(\phi_0^{(II)})^2 - (B+C) = 0. \tag{39}$$

Also we can obtain ρ, P and η for this situation as

$$\begin{aligned} \rho &= V_0 + \frac{\rho_1}{t^{2(\beta\phi_0^{(II)}+1)}}, & (40) \\ P &= \frac{P_1}{t^{2(\beta\phi_0^{(II)}+1)}} + \frac{S}{t}\xi - V_0, & (41) \\ \eta &= \left(\frac{1-S}{2} + \beta\phi_0^{(II)}\right)\frac{t_0^{2\beta\phi_0^{(II)}}}{t^{(2\beta\phi_0^{(II)}+1)}}.l42 & (42) \end{aligned}$$

Here

$$\rho_1 = \left(\frac{S^2-Q}{2} + \left((3\beta^2 - \frac{1}{2})\phi_0^{(II)} - 2\beta S\right)\phi_0^{(II)} - k\right) t_0^{2\beta\phi_0^{(II)}},$$

$$P_1 = \left(\frac{2}{3}S - \frac{Q}{2} - \frac{S^2}{6} + \left(\frac{4}{3}S\beta - 2\beta - (\beta^2 + \frac{1}{2})\phi_0^{(II)}\right)\phi_0^{(II)} - k\right) t_0^{2\beta\phi_0^{(II)}}.$$

So that by using (35) and (5) we can obtain the available $f(R)$ as

$$f(R) = -2\mathcal{V}_0 + F_0 R^{m+1}. \tag{43}$$

Where $F_0 = \frac{w+2k}{w^{m+1}}$ is the constant and $m = \beta\phi_0^{(II)}$,

$$\mathcal{V}_0 = V_0 t_0^{4m}, \tag{44}$$

$$w = 2S - S^2 - Q. \tag{45}$$

For obtain stability of the above $f(R)$, the following condition must be satisfied[17].

$$\frac{d^2 f}{dR^2} > 0, \tag{46}$$

for obtaining this constrain and confined the solution power-low in the expanding universe of Kasner form m should satisfy the following relations[17]

$$m = \beta\phi_0^{II} > 0, \qquad w > -2k. \tag{47}$$

From (43) it is clearly obvious that for $\beta\phi_0^{II} = 0$, the obtained $f(R)$, (43), reduces to $f(R) = R$ which is the standard Einstein general relativity action. However, the quantities ρ, P and η in these two classes for $\beta\phi_0 = 0$ are as

$$\rho = \frac{S^2 - Q}{2t^2}, \tag{48}$$

$$P - \frac{S}{t}\xi = \frac{4S - S^2 - 3Q}{6t^2}, \tag{49}$$

$$\eta = \frac{1 - S}{2t}. \tag{50}$$

The corresponding particular cases are studied in detail in [16]. Considering that it dominant energy condition (DEC) imply that $-\rho + \frac{S}{t}\xi \leq P \leq \rho + \frac{S}{t}\xi$. Then, from (48), (49) and (50), for $P \leq \rho + \frac{S}{t}\xi$ we have that $1 - S \leq 0$ and for of case $-\rho + \frac{S}{t}\xi \leq P$ we obtain that $-2S - S^2 + 3Q \leq 0$. Thus, from these expressions it is easy to see that $\eta \leq 0$ and $\rho \geq 0$.

4.3 CaseIII

In this case we assume $B = -C$, according to (39) $\phi_0^{(III)}$is as

$$\phi_0^{(III)} = \frac{S - 1}{\beta}, \tag{51}$$

so $k = -\frac{C(S-1)(1-6\beta^2)}{2S} = -\frac{\beta(2S-S^2-Q)(S-1)}{2S}$ and the solutions (37- **??**) are as

$$V(t) = V_0 t^{4(S-1)} - k\frac{t^{2(S-2)}}{t_0^{2(S-1)}}, \tag{52}$$

we can obtain physical quantities as

$$\begin{aligned}\rho &= V_0 + \frac{\rho_2}{t^{2S}}, &(53)\\ P &= \frac{P_2}{t^{2S}} + \frac{S}{t}\xi - V_0, &(54)\\ \eta &= \frac{(S-1)}{2}\frac{t_0^{2(S-1)}}{t^{2S-1}}. &(55)\end{aligned}$$

Here

$$\rho_2 = \left(\frac{S^2-Q}{2} - 2S(S-1) + (3\beta^2 - \frac{1}{2})\frac{(S-1)^2}{\beta^2} - k\right) t_0^{2(S-1)}$$

$$P_2 = \left(\frac{2}{3}S - \frac{Q}{2} - \frac{S^2}{6} + \frac{4}{3}S(S-1) - 2(S-1) - (\beta^2 + \frac{1}{2})\frac{(S-1)^2}{\beta^2} - k\right) t_0^{2(S-1)}$$

and we can obtain $f(R)$ as

$$f(R) = -2\mathcal{V}_0 + F_0 R^{S-1}. \qquad (56)$$

Whereas the shear coefficient of viscosity at early time is much bigger than the bulk viscosity [18], so we take $\xi = 0$. And also according to (55) we can assume that V_0 will be equal to Λ, therefore we can neglect it with respect to other terms in (53), (54). Hence in this case one can write the barotropic equation of state as $P = \gamma\rho$. It is well known that the dominant energy condition (DEC) imply that γ lies in the interval [-1, 1]. From (53) and (54), and the above assumptions, we can obtain the barotropic equation of state as

$$\gamma = \frac{-k + \frac{2}{3}S - \frac{Q}{2} - \frac{S^2}{6} + \frac{4}{3}S(S-1) - 2(S-1) - (\beta^2 + \frac{1}{2})\frac{(S-1)^2}{\beta^2}}{-k + \frac{S^2-Q}{2} - 2S(S-1) + (3\beta^2 - \frac{1}{2})\frac{(S-1)^2}{\beta^2}}. \qquad (57)$$

The quantity γ has four independent parameters $\beta, p_i s(1 = 1, 2, 3)$. We want to obtain the available range of these parameters in which the dominant energy condition, $-1 \leq \gamma \leq 1$ and $S > 1$ is satisfied. We have drown the equation of state parameter, γ versus p_3 for two set of parameters. *a*) for $\beta = \frac{1}{\sqrt{6}} = 0.41, p_1 = 0.5$ and $p_2 = 0.3$. *b*) for $\beta = 0.5, p_1 = 0.5$ and $p_2 = 0.3$ in figure 1. For these two considerations we obtain two different intervals for p_3 as (0.424 , 0.69) and (0.456 , 0.69) respectively. It is seen that these two set of parameters and corresponding intervals of p_3 are satisfying the constraint $S > 1$ and dominant energy condition. Also, according to[6] the second law of thermodynamics imply that $\eta > 0$ hence, considering to equation (55), it is clearly seen that this condition is satisfied. Therefore we derived that this model satisfy the dominant energy condition (DEC) and the second law of thermodynamics (SLT), simultaneously. From these two sets of data seen that by increasing the value of β, the constraint $S > 1$, DEC and SLT is satisfied for smaller interval of p_3.i.e $\triangle p_3^{(a)} = 0.266 > \triangle p_3^{(b)} = 0.234$.



(a) (b)

Figure 1: Variation the state parameter, γ with respect to p_3: (a); In this subfigure we draw γ for $p_1 = 0.5$, $p_2 = 0.3$ and $\beta = \frac{1}{\sqrt{6}} = 0.41$. We have obtained the interval $(0.424, 0.69)$ for p_3 in which the dominant energy condition is satisfied. (b);In this subfigure we draw γ for $p_1 = 0.5$, $p_2 = 0.3$ and $\beta = 0.5$. We have obtained the interval $(0.456, 0.69)$ for p_3 in which the dominant energy condition is satisfied.

5 Conclusion

The main purpose of the present work is to study of the $f(R)$ theory of gravity in an anisotropic metric, such as Kasner form metric. We have assumed that the universe is filled by a viscose cosmic fluid which is endowed with a shear viscosity η and bulk viscosity ξ. In this work, we have obtained the functions of $f(R)$ gravity that satisfied condition stability and expanding universe of Kasner form metric. We derived that this model satisfy the dominant energy condition (DEC) and the second law of thermodynamics (SLT), simultaneously. At last, by plotting the equation of state pamperer, γ, versus one of the Kasner's index , p_3, for two set of parameters, we have obtained two intervals for p_3 in which the constraint $S > 1$, DEC and SLT are satisfied. Therefore, it is clear that $f(R)$ gravity model as well as satisfied solutions, viscous cosmology in scalar theories for Kasner type metrics.

References

[1] A. G. Riess *et al.*, L. **116**, 1009 (1998); S. Perlmutter *et al.*, Astrophys. J. **517**, 565 (1999).

[2] D. N. Spergel *et al.*, Astrophys. J. suppl.**148**, 175 (2003).

[3] M. Tegmark *et al.*, phys.Rev. **D 69**, 103501 (2004).

[4] R. M. Wald., General Relativity, (Chicoago and London: The University of Chicago Press) (1984).

[5] S. Weinberge., Gravitational and Cosmology , (New York: Willey) (1984).

[6] C. W. Misner, K. S. Thorne and J. A. Wheeler., Gra-vitation (San Francisco: W. H. Freeman and company) (1970).

[7] Bijan Saha, Phys. Rev. D **64**, 123501 (2001) [arXiv: gr-qc/0107013].

[8] Bijan Saha and Todor Boyadjiev, Phys. Rev. D **69**, 124010 (2004) [arXiv: gr-qc/0311045].

[9] I. M. Khalatnikov and A.Yu. kamenshchik, Phys. Lett. B **553**, 119 (2003).

[10] E. F. Bunn, P. G. Ferreira, and J. Silk, Phys. Rev. Lett. **77**, 2883 (1996).

[11] E. Bertschinger, Physica D **77**, 354 (1994).

[12] E. Guzman, Astrophys. Space Sci. **199**, 289 (1993).

[13] P. Halpern, Gen. Relativ. Gravit. **26**, 781 (1994).

[14] R. J. van den Hoogen, Class. Quantum Grav. **12**, 2335 (1995).

[15] D. Kalligas, P. S. Wesson, and C. W. F. Everitt, Gen. Relativ. Gravit. **27**, 645 (1995).

[16] I. Brevik, S.V. Pettersen, Phys. Rev. D **56**, 3322 (1997).

[17] A. Aghmohammadi, Kh. Saaidi, M. R. Abolhassani, A. Vajdi, Phys. Scripta **80**, 065008 (2009).

[18] N. Caderni, R. Fabbri, Phys. Lett. B **69**, 508 (1977).